



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The Ubiquitin-proteasome System as a Regulator of Plant Immunity

Citation for published version:

Adams, EHG & Spoel, SH 2018, 'The Ubiquitin-proteasome System as a Regulator of Plant Immunity', *Journal of Experimental Botany*. <https://doi.org/10.1093/jxb/ery216>

Digital Object Identifier (DOI):

[10.1093/jxb/ery216](https://doi.org/10.1093/jxb/ery216)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Experimental Botany

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The ubiquitin-proteasome system as a transcriptional regulator of plant immunity

Eleanor H. G. Adams and Steven H. Spoel

Institute of Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh,
Edinburgh EH9 3BF, United Kingdom

Corresponding author:

Dr. Steven H. Spoel (ORCID: [0000-0003-4340-7591](https://orcid.org/0000-0003-4340-7591))

Institute of Molecular Plant Sciences

School of Biological Sciences

University of Edinburgh

King's Buildings

Max Born Crescent

Edinburgh, EH9 3BF

United Kingdom

Email: steven.spoel@ed.ac.uk

Phone: +44 (0)131 650 7065

Date of submission: 14 Feb. 2018

Number of figures: 1

Word count: 4,850

Running title: UPS-dependent immune transcription

Keywords: Ubiquitin proteasome system; Cullin-RING; NPR1; WRKY transcription factors; salicylic acid; plant immunity

Highlights

Activation of plant immunity is associated with dramatic changes in gene expression. Here we discuss diverse roles of the ubiquitin-proteasome system as a transcriptional regulator of immune genes.

Abstract

The ubiquitin-proteasome system (UPS) has been shown to play vital roles in diverse plant developmental and stress responses. The UPS post-translationally modifies cellular proteins with the small molecule ubiquitin, resulting in their regulated degradation by the proteasome. Of particular importance is the role of the UPS in regulating hormone-responsive gene expression profiles, including those triggered by the immune hormone salicylic acid (SA). SA utilises components of the UPS pathway to reprogram the transcriptome for establishment of local and systemic immunity. Emerging evidence has shown that SA induces the activity of Cullin-RING ligases (CRLs) that fuse chains of ubiquitin to downstream transcriptional regulators and consequently target them for degradation by the proteasome. Here we review how CRL-mediated degradation of transcriptional regulators may control SA-responsive immune gene expression programmes and discuss how the UPS can be modulated by both endogenous and foreign exogenous signals. The highlighted research findings paint a clear picture of the UPS as a central hub for immune activation as well as a battle ground for hijacking by pathogens.

Introduction to the Ubiquitin-Proteasome System

Regulated degradation of short-lived or damaged proteins plays vital roles in the cellular development and signalling across eukaryotes. The ubiquitin-proteasome system (UPS) is responsible for the selection, targeting and proteolysis of specific substrates destined for degradation. UPS components are especially abundant in plants, suggesting this system is a key hub for the regulation of numerous plant cell processes. Emerging evidence from plants and animals indicates that UPS components do not only function merely to target substrates for degradation; they are also critical transcriptional co-regulators that are indispensable for developmental and stress-responsive gene expression programmes (Geng *et al.*, 2012; Kelley and Estelle, 2012; Santner and Estelle, 2009; Vierstra, 2009).

Central to the functioning of the UPS is the post-translational modification of substrates by a single or polymeric chain of ubiquitin, a highly conserved small 8.5 kDa protein. Ubiquitin is covalently added to lysine (Lys) residues of substrates in a multi-step enzymatic cascade that involves E1 activating, E2 conjugating and E3 ligase enzymes. First an E1 enzyme forms a high-energy thioester bond to an ubiquitin adduct, which is then transferred onto the active site cysteine (Cys) residue of an E2 enzyme. The ubiquitin-loaded E2 enzyme then partners with an E3 ligase to transfer ubiquitin to a Lys residue of the target substrate. Reiterations of this reaction allow subsequent ubiquitin molecules to be similarly attached to internal Lys residues of the preceding ubiquitin, thus generating a chain of polyubiquitin on the substrate (Komander and Rape, 2012; Smalle and Vierstra, 2004). While chains can form by linking different Lys residues of ubiquitin, Lys48 linkage between four or more ubiquitins exhibits high affinity for ubiquitin receptors of the proteasome, a large 2.5 MDa ATP-dependent chambered protease consisting of dozens of distinct subunits (Pickart and Cohen, 2004; Thrower *et al.*, 2000). The 19S regulatory cap of the proteasome is responsible for recognition of ubiquitinated substrates, the chaperone-assisted unfolding of substrates, and releasing polyubiquitin for recycling. Subsequently unfolded substrates are threaded into the 20S particle of the proteasome, a barrel-shaped multi-catalytic proteinase, where they are cleaved into peptides (Pickart and Cohen, 2004).

Compared to other eukaryotes, plant genomes often encode for large numbers of UPS components, suggesting the UPS plays important roles in diverse cellular processes. Recent years have clearly shown that the UPS contributes to the establishment of local and systemic immunity in plants. Comprehensive reviews on the role of ubiquitination in plant immunity are already available (Duplan and Rivas, 2014; Furniss and Spoel, 2015; Marino *et al.*, 2012; Trujillo and Shirasu, 2010), so here we provide a more focussed update on our understanding of how selected components of the UPS function as transcriptional co-regulators of plant immune genes.

SA-responsive Cullin-RING Ligases are transcriptional co-regulators

Amongst UPS components, E3 ligases are predominantly responsible for establishing selective UPS activity. E3 ligases specifically recognise and interact with their substrates, leading to their (poly)ubiquitination and, in case of Lys48-linked ubiquitin chains, subsequent degradation by the proteasome. Plants contain large numbers of E3 ligases (e.g. the *Arabidopsis* genome encodes for >1,500) that are predicted to each target different substrate repertoires (Mazzucotelli *et al.*, 2006). Among these the family of modular multi-subunit Cullin-RING Ligases (CRL), which are predicted to form nearly 700 different E3 ligases, have been shown to be necessary for plant defence signalling by the immune hormones salicylic acid (SA) and jasmonic acid (JA).

Biotropic pathogen attack leads to rapid accumulation of SA, which acts as both a local and systemic signal for the induction of appropriate defences (Spoel and Dong, 2012). In incompatible plant-pathogen interactions SA acts locally as an agonist of programmed cell death, which is thought to confine pathogens to a hostile environment and deprive them of further nutrients. Pathogen attack also leads to accumulation of SA in tissues adjacent or distant from the (attempted) infection site where it coordinates the reprogramming of ~2,200 genes, including *pathogenesis-related (PR)* genes (Wang *et al.*, 2006). Genetic screens for SA-insensitive *Arabidopsis* mutants have repeatedly identified different *npr1* (*non-expressor of PR genes 1*) alleles (Cao *et al.*, 1994; Delaney *et al.*, 1995; Glazebrook *et al.*, 1996; Shah *et al.*, 1997). NPR1 encodes a transcriptional coactivator with two protein-protein interaction domains: an N-terminal BTB (Bric-à-brac Tramtrack, and Broad complex) domain and a more C-terminal Ankyrin repeat domain (Cao *et al.*, 1997; Ryals *et al.*, 1997; Tada *et al.*, 2008). In resting cells conserved Cys residues in and adjacent to the BTB domain form disulphide bonds, resulting in formation of a high molecular weight cytoplasmic NPR1 oligomer that is excluded from the nucleus (Kinkema *et al.*, 2000; Mou *et al.*, 2003). Accumulation of SA triggers changes in cellular redox potential that together with the thiol reductase action of Thioredoxins lead to reduction of these disulphide bonds with subsequent release of NPR1 monomer that translocates to the nucleus (Kinkema *et al.*, 2000; Mou *et al.*, 2003; Tada *et al.*, 2008). Nuclear NPR1 monomer interacts with and trans-activates transcription factors of the TGA and WRKY families that associate with SA-responsive gene promoters (Boyle *et al.*, 2009; Després *et al.*, 2000; Saleh *et al.*, 2015; Zhang *et al.*, 1999; Zhou *et al.*, 2000). Thus, NPR1 is thought to be a master coactivator of SA-responsive immune gene transcription.

Disrupting formation of the NPR1 oligomer by mutation of oxidant-sensitive Cys156 led to loss of long-term SA-induced resistance, indicating the oligomer is indispensable for NPR1 homeostasis (Tada *et al.*, 2008). This effect was associated with a drastic decrease in NPR1 protein levels, suggesting nuclear NPR1 protein is unstable. Indeed, pharmacological

inhibition of the proteasome resulted in accumulation of NPR1 in the nucleus of resting cells and constitutive activation of its direct target genes (Spoel *et al.*, 2009). Co-immunoprecipitation experiments demonstrated that NPR1 associated with a nuclear CRL3 ligase (also known as BC3B for BTB/Cullin3/BTB). Genetic perturbation of CRL3 by mutation of the Cullin 3 subunit or by mutating the COP9 signalosome that regulates the stability and activity of Cullin proteins (Petroski and Deshaies, 2005), stabilised NPR1 protein and was associated with constitutive expression of its target genes. This suggests that CRL3-mediated degradation of NPR1 functions to maintain SA-responsive immune gene expression in a latent state, thereby preventing onset of autoimmunity (Figure 1).

Despite being necessary for the SA-responsive activation of target genes, paradoxically SA-induced NPR1 was also a substrate for CRL3-mediated ubiquitination and proteasomal degradation (Spoel *et al.*, 2009). Mutation of CRL3 or an NPR1 phospho-site responsible for recruitment of NPR1 to CRL3 led to reduced SA-responsive gene expression and impaired disease resistance. These findings suggest that instability of NPR1 appeared to be necessary for full activation of its target genes. As we have proposed previously (Furniss and Spoel, 2015; Skelly *et al.*, 2016; Spoel *et al.*, 2009), this may be due to gene expression requiring continuous delivery of fresh transcriptionally competent NPR1 to active gene promoters (Figure 1). Although most transcriptional regulators are stable proteins, selected eukaryotic transcriptional regulators have been found to exhibit a similar instability as NPR1. These regulators often contain overlapping sequences that act as transactivation domains and degradation motifs (Salghetti *et al.*, 2000). NPR1 was found to form a trans-activating transcriptional complex with TGA2 transcription factors, which required core residues of the BTB domain as well as oxidation of two C-terminal cysteine residues (Rochon *et al.*, 2006). Although these transactivation domains differ from the N-terminal phosphorylation sites that are necessary for recruitment of NPR1 to CRL3 (Spoel *et al.*, 2009), it is plausible that Lys ubiquitination occurs in or near these domains.

In rice, the *Oryza sativa* WRKY45 transcriptional activator exhibits overlap between trans-activating and proteolysis targeting sequences. OsWRKY45 performs a very similar function as Arabidopsis NPR1, as it is responsible for SA-responsive transcriptional reprogramming and establishment of resistance against bacterial and fungal pathogens (Nakayama *et al.*, 2013; Shimono *et al.*, 2012; Shimono *et al.*, 2007). Pharmacological treatment with proteasome inhibitors resulted in accumulation of ubiquitinated OsWRKY45 and blocked SA-induced transcriptional activation of its target genes (Matsushita *et al.*, 2013). Importantly, C-terminal sequences in OsWRKY45 were necessary for both its transcriptional activity as well as UPS-dependent degradation. Thus, SA-responsive gene expression in higher plants may be dependent on transcriptional regulators that harbour

sequences that function as both trans-activating domains and UPS targeting signals (Figure 1).

While it remains unknown which E3 ligase is responsible for targeting OsWRKY45 for proteasome-dependent degradation, a CRL3 ligase has been implicated in SA-dependent immunity in rice. Studies on *Cullin 3a* (*OsCUL3a*) revealed that it interacted with RING-BOX1 (RBX1) and RBX2 to constitute the core of CRL3 (Liu *et al.*, 2017). Genetic analyses showed that *oscul3a* mutants displayed typical symptoms of autoimmune activation, suggesting CRL3 functions as immunosuppressant in rice. Similar to Arabidopsis NPR1, OsWRKY45 is continuously degraded by the UPS in resting cells and failure to clear this activator results in autoimmune phenotypes reminiscent of *oscul3a* (Matsushita *et al.*, 2013). Thus, it is plausible that in analogy to Arabidopsis, CRL3 also targets OsWRKY45 for proteasome-mediated degradation in rice (Figure 1).

OsCUL3 was found to physically associate with the rice homologue of NPR1, known as OsNPR1 or NH1, which also functions in an SA-responsive immune signalling pathway. Cycloheximide and proteasome inhibition assays established that OsCUL3 is necessary for the proteasome-dependent degradation of OsNPR1, indicating that OsNPR1 is also a substrate of CRL3 (Liu *et al.*, 2017). Rice CRL3 has the potential to influence a large transcriptional immune programme, as accumulation of OsNPR1 protein in *oscul3a* mutants was associated with activation of *PR* genes. This is in agreement with previous reports demonstrating that overexpression of OsNPR1 resulted in constitutive activation of immune genes and resistance to bacterial blight (Chern *et al.*, 2001; Chern *et al.*, 2005; Yuan *et al.*, 2007). However, genome-wide transcript profiling of *OsNPR1* knock down lines showed that its function as an activator of immune genes was relatively modest in comparison to its role in transcriptional suppression. OsNPR1 directly or indirectly down regulated the expression of genes involved in photosynthesis and in chloroplast translation and transcription, suggesting it plays an important role in resource reallocation during establishment of immunity (Sugano *et al.*, 2010). How CRL3-mediated ubiquitination of OsNPR1 affects suppression of these growth and development related genes currently remains unknown. Considering CRL3 controls the cellular levels of OsNPR1, it is expected that CRL3 plays an important role in limiting the suppressive effects of NPR1, thereby managing the balance between defence and growth (Figure 1).

Composition and potential structure of CRL3 ligases in SA signalling

To recruit substrates CRL3 ligases utilize specific substrate adaptors that typically contain a BTB domain and one or more additional protein-protein interaction domains. The BTB domain and adjacent sequences directly interact with CUL3 (Canning *et al.*, 2013; Stogios *et al.*, 2005; Zhuang *et al.*, 2009), while the additional protein-protein interaction domain binds

to the substrate. Intriguingly, the *NPR* protein family appears to have all the necessary features to function as substrate adaptors for CUL3. In *Arabidopsis* this family consists of *NPR1* and five *NPR1*-like genes, namely *NPR1-like 2* (*NPR2*), *NPR3*, *NPR4*, *BLADE-ON-PETIOLE2* (*BOP2*; also named *NPR5*), and *BOP1* (also named *NPR6*). Each of the *NPR1* proteins contains a conserved BTB and Ankyrin-repeat domain. Whereas *BOP1* and *BOP2* contribute to leaf development and JA-mediated immune signalling (Canet *et al.*, 2012; Hepworth *et al.*, 2005), *NPR1*–*4* have all been implicated in SA-dependent immune signalling (Canet *et al.*, 2010; Cao *et al.*, 1997; Fu *et al.*, 2012; Liu *et al.*, 2005; Zhang *et al.*, 2006). While little is known about the role of *NPR2* in defence signalling, it was demonstrated that genetically *NPR2* contributes to SA perception in *npr1* null mutants (Canet *et al.* 2010). Recent work indicated that recruitment of *NPR1* to CUL3 was dependent on both *NPR3* and *NPR4*, suggesting these proteins recruit *NPR1* for ubiquitination by a CUL3^{*NPR3/4*} ligase. This ligase was found to be necessary for establishment of SA-dependent systemic acquired resistance and pathogen effector-induced programmed cell death responses (Fu *et al.*, 2012). Although it remains unclear how and if *NPR3* and *NPR4* regulate the transcriptional activity of *NPR1*, these results clearly indicate the potential for *NPR* protein family members to provide specificity to CUL3 ligases in plant immunity.

Computational and protein crystallisation data have shown that CUL3 ligases are dimeric (Stogios *et al.*, 2007; Zhuang *et al.*, 2009; Zimmerman *et al.*, 2010). Dimerization is driven by tight electrostatic interactions between BTB domains of two substrate adaptors, allowing the binding of two CUL3 subunits. Rather than recruiting two substrates, CUL3 dimerisation has been suggested to improve the efficiency by which a single substrate molecule is ubiquitinated. It is thought that the dimer constrains the mobility of the substrate, thereby improving the rate of ubiquitination on target lysines (McMahon *et al.*, 2006). Because self-association was found to be a general feature of many CUL3 ligases (McMahon *et al.*, 2006), it is likely that *NPR3* and *NPR4* allow CUL3 in plants to form a dimeric complex that recruits *NPR1* for ubiquitination. Current work in our laboratory is investigating if CUL3 forms homo- or heterodimers. Although heterodimers are not documented as a feature of BTB-containing CUL3 substrate adaptors, it has been demonstrated for two hetero-oligomerising F-box proteins, Pop1p and Pop2p, which are part of an Skip-Cullin-F-box (SCF, also known as CUL1) ligase in fission yeast (Seibert *et al.*, 2002). While heterodimeric SCF^{Pop1p-Pop2p} target the cyclin-dependent kinase inhibitor Rum1p for degradation, homodimeric SCF^{Pop1p} and SCF^{Pop2p} complexes probably have different substrate preferences. Thus heterodimer or even heterooligomer formation between different *NPR* proteins potentially increases combinatorial diversity in substrate preference that could extend well beyond *NPR1* as the sole substrate.

Coactivator turnover is modulated by endogenous and exogenous signals

The destructive nature of the UPS must be tightly controlled to ensure appropriate levels of substrate degradation. A major regulatory checkpoint is the selective recruitment of substrates to E3 ligases. Diverse cellular signals including post-translational modifications mark substrates for recruitment to E3 ligases. Activation of nuclear NPR1 and its recruitment to CRL3 was recently shown to involve complex interplay between SUMOylation and (de)phosphorylation (Saleh *et al.*, 2015). SA-induced dephosphorylation of Ser55/59 was prerequisite for modification of NPR1 by SUMO3. NPR1 SUMOylation was proposed to regulate positional interactions with its target promoters through differential association with transcription factor partners. Whereas unmodified NPR1 interacted with the transcriptional repressor WRKY70, SUMOylated NPR1 preferentially associated with the transcriptional activator TGA3. Importantly, SUMOylation itself or the resulting switch in transcriptional partner (*i.e.* from WRKY70 to TGA3) was required for subsequent phosphorylation of Ser11/15 (Saleh *et al.*, 2015). Ser11/15 phosphorylation was in turn necessary for recruitment of NPR1 to CRL3, resulting in its ubiquitination and turnover by the proteasome (Figure 1) (Spoel *et al.*, 2009). Although the exact residues remain unknown, OsWRKY45 in rice was also found to be phosphorylated. Interestingly, the phosphorylated form was highly responsive to proteasome inhibition, suggesting that site-specific phosphorylation of OsWRKY45 may also be required for its UPS-dependent degradation (Figure 1) (Matsushita *et al.*, 2013). Thus, extensive interplay between diverse post-translational signals regulates the stability and associated activity of SA-responsive transcriptional (co)activators (Skelly *et al.*, 2016).

Recruitment of substrates to E3 ligases is not only regulated by post-translational control mechanisms, it may also be facilitated or inhibited by small molecules. While this has driven the design of synthetic molecules for human medicine (Zheng and Shabek, 2017), in plants several major developmental and stress signalling pathways naturally utilise E3 ligases as receptors for small-molecule hormones. This was first discovered for the plant developmental hormone auxin, which promotes the recruitment of a family of transcriptional repressors to an SCF^{TIR1} ubiquitin ligase. Structural biology approaches have shown that auxin acts as ‘molecular glue’ by enhancing protein-protein interactions between the SCF F-box subunit TIR1 and transcriptional repressors. Consequently, auxin perception at or near the chromatin relieves transcriptional suppression by SCF^{TIR1}-mediated ubiquitination and degradation of repressors (Dharmasiri *et al.*, 2005; Kepinski and Leyser, 2005; Santner and Estelle, 2009). Other hormones, such as JA-isoleucine, are perceived similarly by different SCF ligase–substrate complexes, indicating that hormone perception by E3 ligases is a key mechanism for direct transcriptional regulation (Kelley and Estelle, 2012). Intriguingly, SA is also perceived by the CRL3 substrate adaptors NPR3 and NPR4, suggesting hormone

perception may be common a mechanism of perception by CRL ubiquitin ligases. Distinct from other hormones, however, SA also regulates how and when NPR3 and NPR4 interact with NPR1. This is a consequence of the vastly different affinities NPR3 and NPR4 have for SA. Whereas NPR3 has a relatively low affinity for SA, NPR4 displays much higher affinity. Moreover, SA binding has opposing effects on substrate recruitment. The result is that elevated levels of SA disrupt NPR1-NPR4 interactions, but promote NPR1-NPR3 interactions (Fu *et al.*, 2012). Thus, in absence of pathogen threat when low levels of SA are present, the CUL3^{NPR4} ubiquitin ligase complex is thought to suppress NPR1 monomer levels to prevent autoimmunity. Conversely, when cellular levels of SA increase following pathogen attack, NPR1 recruitment is switched from CUL3^{NPR4} to CUL3^{NPR3}, which is necessary for local and systemic immune responses (Fu *et al.*, 2012; Furniss and Spoel, 2015). Although details of the mechanisms by which SA regulates CUL3^{NPR3/NPR4} remain largely unknown, the clear analogy to other CRL-dependent signalling pathways demonstrates that SA is a key small molecule involved directly in the regulation of NPR1 stability and therefore likely also NPR1 transcriptional potency.

Endogenous signals originating from hormone signalling pathways may also modulate CUL3-mediated ubiquitination and degradation of NPR1 coactivator. The developmental hormone abscisic acid (ABA) interacts antagonistically with SA-dependent immune signalling. Some pathogens hijack this antagonism by inducing apparent increases in ABA biosynthesis, thereby inhibiting signalling steps both up and downstream of SA accumulation (de Torres-Zabala *et al.*, 2007; Mohr and Cahill, 2007; Yasuda *et al.*, 2008). A recent report investigated how ABA might impact SA signalling by examining NPR1 stability (Ding *et al.*, 2016). It was found that treatment with ABA strongly reduced the cellular level of NPR1 in a CUL3^{NPR3/NPR4} and proteasome-dependent manner, suggesting that in unchallenged cells ABA antagonised SA signalling by destabilising NPR1. However, by changing the timing of pharmacological applications of SA and ABA, it was found that ABA reduces NPR1 protein levels only if ABA treatment preceded SA treatment. These data suggest that ABA has less control over the stability of SA-induced NPR1. Indeed, phosphorylation of Ser11/15, which is necessary for SA-induced NPR1 degradation, appeared to block ABA-induced NPR1 instability. Nonetheless, ABA treatment could strongly reduce SA-induced *PR-1* gene expression, implying that antagonisms between SA and ABA is more complex than can be observed by examining NPR1 protein levels at a single time point. By temporally surveying NPR1 protein levels during infection by the virulent bacterial pathogen *Pseudomonas syringae*, the authors revealed that SA and ABA accumulate sequentially, which may allow a switch from SA-induced NPR1 protein degradation to ABA-induced turnover (Ding *et al.*, 2016). The functional outcome of switching between mechanisms of NPR1 degradation remains unclear but ABA-induced degradation of NPR1

during later stages of the immune response might be necessary for full-scale induction of NPR1-dependent target genes. How ABA recruits NPR1 for degradation by CRL3^{NPR3/4} also remains unknown, but it may well involve phosphorylation of NPR1 by members of the SNF1-related protein kinase (SnRK) family. SnRK members have been implicated in ABA signalling by promoting the transcriptional activity of ABA-responsive coactivators. More recently, SnRK2.8 was found to phosphorylate NPR1 at Ser589 and possibly Thr373, allowing its nuclear translocation in tissues distal from the site of infection (Lee *et al.*, 2015). It is plausible that coordination between ABA- and SA-induced phosphorylation events orchestrate diverse pathways of NPR1 ubiquitination and degradation, each with distinct transcriptional outputs.

In addition to endogenous inputs, exogenous signals may also modulate transcriptional coactivator turnover. Curiously, pathogen effectors from a variety of plant pathogens have been shown to interfere with components of the host UPS machinery. Some effectors have been found to inhibit the activity of immune-related E3 ligases or enhance their stability, while others such as *P. syringae* avrPtoB mimic RING- and U-box-type E3 ligases and target host pathogen recognition receptors for degradation (Duplan and Rivas, 2014). More recently it was shown that the infection strategy of *P. syringae* includes inhibition of host proteasome activity in a type III secretion-dependent manner (Üstün *et al.*, 2016). A screen for secreted effectors uncovered several proteins, including HopM1, with proteasome inhibitor activity. Co-immunoprecipitation experiments showed that HopM1 complexed with a variety of E3 ligases and proteasome subunits, demonstrating it directly targets the UPS (Üstün *et al.*, 2016). Other pathogen effectors have also been identified to directly target the proteasome. The *Xanthomonas campestris* effector protein XopJ was found to suppress host proteasome activity by degrading the proteasomal AAA-ATPase subunit RPT6 (Üstün *et al.*, 2013; Üstün and Bornke, 2015). The ATPase activity of RPT6 is thought to be required for the unfolding of substrates prior to their insertion into the 20S catalytic barrel. RPT6 contains Walker A and Walker B motifs that are essential for its ability to bind and hydrolyse ATP, respectively. Interestingly, it was shown that mutation of the Walker A motif abolished interaction with XopJ, whereas mutation of the Walker B motif prevented XopJ-mediated proteolysis of RPT6 (Üstün and Bornke, 2015). These findings suggest that only ATP-bound RPT6 is recognised by XopJ and that XopJ may mimic host substrates intended for proteasomal degradation. Importantly, XopJ-mediated proteolysis of RPT6 was linked to increased accumulation of ubiquitinated NPR1 and decreased turnover of this transcriptional coactivator, likely preventing full-scale activation of SA-responsive immune genes (Spoel *et al.*, 2009; Üstün and Bornke, 2015).

Because of its critical role in the activation of SA-responsive immune genes, it has long been speculated that pathogen effectors also directly target NPR1 and suppress its

transcriptional coactivator activity. Indeed, recent work identified the *P. syringae* effector avrPtoB, a U-box type E3 ligase, as an interactor of NPR1 (Chen *et al.*, 2017). Curiously, SA enhanced interaction between avrPtoB and NPR1, leading to NPR1 ubiquitination and subsequent proteasomal degradation. Unlike immune-induced NPR1 ubiquitination and turnover, avrPtoB-induced degradation of NPR1 negatively affected SA-responsive gene expression and immunity (Chen *et al.*, 2017). This suggests that avrPtoB either ubiquitinates distinct Lys residues in NPR1 as compared to CRL3 or it targets NPR1 for ubiquitination prior to its recruitment to CRL3. As SA promotes interaction between avrPtoB and NPR1, it is plausible that avrPtoB titrates NPR1 away from SA-mediated binding to the CRL3 substrate adaptor NPR3.

Degradation of other SA-responsive transcriptional immune regulators

The UPS may control SA-responsive gene expression in ways that go well beyond regulating the stability of master (co)activators such as NPR1 and OsWRKY45. SA-responsive gene expression is modulated by many other transcriptional activators and repressors from the TGA, WRKY and NIMIN or OsNRR families, many of which physically interact with NPR1 and the CRL3 substrate adaptors NPR3 and NPR4 (Chern *et al.*, 2014; Liu *et al.*, 2005; Saleh *et al.*, 2015; Shi *et al.*, 2013; Weigel *et al.*, 2001; Zhang *et al.*, 2006). Interestingly, some of these transcriptional regulators or their close relatives have been reported to exhibit UPS-dependent instability. TGA1 and TGA3 transcriptional activators have been shown to interact with NPR proteins and their protein levels are regulated by post-transcriptional mechanisms. Although their protein levels appeared to be controlled developmentally, inhibition of the proteasome resulted in accumulation of TGA3 in the nucleus (Pontier *et al.*, 2002). This suggests that TGA3 is unstable and its degradation could be managed by SA. Because TGA3 interacts with NPR3 and NPR4 (Zhang *et al.*, 2006), it is plausible that TGA3 is regulated by an SA-induced CRL3^{NPR3} or CRL3^{NPR4}. Alternatively, TGA3 could be targeted for degradation indirectly through its interaction with NPR1 via concurrent ubiquitination and degradation. Concurrent ubiquitination and degradation of multiple physically associated substrates has already been reported for a CRL3 ligase involved in light signalling (Ni *et al.*, 2014). In addition to TGA transcription factors, WRKY transcription factors regulate SA-responsive gene expression both positively and negatively. While (in)stability of the wider WRKY protein family has not yet been examined, some WRKY proteins such as the above discussed OsWRKY45 have been found to be subject to UPS-dependent degradation. In Arabidopsis WRKY53 was found to be a substrate of HECT domain-containing Ubiquitin Protein Ligase 5 (UPL5) during leaf senescence (Miao and Zentgraf, 2010). Notably WRKY53 is also an activator of SA-responsive immune genes (Wang *et al.*, 2006), implying the possibility that control of WRKY53 protein levels by the

UPS also impacts SA-dependent gene expression. In Chinese wild grapevine *Vitis pseudoreticulata* WRKY11 (VpWRKY11) was targeted for ubiquitination and degradation by the RING E3 ligase *Erysiphe necator*-induced RING finger protein 1 (EIRP1), which was necessary for resistance to a variety of different pathogens but specific effects on SA-responsive gene expression remained unclear. (Yu *et al.*, 2013). Finally, in *Arabidopsis* Signal Responsive 1 (SR1), a Ca²⁺/calmodulin binding transcription factor, was found to be controlled by ubiquitination and degradation. SR1 binds to and suppresses the promoter of *Enhanced Disease Susceptibility 1 (EDS1)*, a gene involved in the biosynthesis of SA. Consequently, mutant *sr1-1* plants exhibit increased transcript levels of *EDS1* as well as other SA biosynthesis genes and accumulate elevated levels of SA (Du *et al.*, 2009). Recent work demonstrated that SR1 is recruited to a CRL3 ligase for ubiquitination and proteasome-mediated degradation. Interestingly, SR1 is recruited to CRL3 by SR1 Interacting Protein 1 (SR1IP1), a protein containing both BTB and non-phototropic hypocotyl 3 (NPH3) protein-protein interaction domains, which are typical characteristics of a CRL3 adaptor (Stogios *et al.*, 2005; Zhang *et al.*, 2014). SR1IP1 was shown to function as a positive regulator of SA-mediated defence responses by removing the transcriptional repressor SR1. Taken together with knowledge gained on CRL3^{NPR} ligases, these findings imply the exciting possibility that CRL3 dynamically switches between different substrate adaptors to recruit distinct transcriptional (co)regulators for ubiquitination and degradation.

Perspectives

The complex roles of the UPS in regulating eukaryotic gene expression have been an intense field of study for some time now (Collins and Tansey, 2006; Geng *et al.*, 2012). Involvement of the UPS in plant immune transcriptional reprogramming is now well established but the complexity is only just being uncovered. Aside from regulating SA-mediated immunity, the UPS is also vital in the control of JA-responsive gene expression during development and immunity. JA facilitates the molecular association between SCF^{COI1} ligase and its substrates, JAZ transcriptional corepressors. SCF^{COI1}-mediated degradation of JAZ corepressors releases numerous transcription factors from suppression and leads to the activation of amongst others defence responses against necrotrophic pathogens and insects, a topic extensively reviewed elsewhere (Goossens *et al.*, 2016; Zhang *et al.*, 2017). Interestingly, under many circumstances the SA and JA signals are antagonistic. SA exerts its antagonisms through the function of NPR1, which was uncovered as a potent suppressor of JA-responsive gene expression (Spoel *et al.*, 2003). It remains unclear if there are any spatial chromatin interactions between SA-responsive CRL3^{NPR} and JA-responsive SCF^{COI1} ligases but evidence suggests that SA and NPR1 suppress JA signalling further downstream. Indeed, activation of SA signalling failed to interfere with SCF^{COI1}-mediated

degradation of JAZ corepressors. Instead, SA strongly reduced protein levels of the JA-responsive transcriptional activator ORA59, which functions downstream of SCF^{COI1} (Van der Does *et al.*, 2013). Future research should reveal if this negative effect of SA on ORA59 protein levels is mediated by an SA-induced CRL3^{NPR} ligase. This is highly plausible as a recent report demonstrated that CRL3^{NPR3/NPR4} may also target transcriptional components of JA signalling during pathogen effector-triggered immunity (Liu *et al.*, 2016). Unlike local immune responses to virulent pathogens, immunity triggered by the recognition of pathogen effectors is not associated with antagonisms between the SA and JA signals (Spoel *et al.*, 2007). It was found that antagonism is avoided through CRL3^{NPR3/NPR4}-mediated ubiquitination and removal of JAZ corepressors, allowing activation of JA-responsive genes in a cellular environment of active SA signalling. Curiously, pharmacological application of SA was insufficient to induce degradation of JAZ corepressors by CRL3^{NPR3/NPR4}, suggesting that effector recognition triggers additional signalling pathways that activate or recruit this E3 ligase. Taken together, these findings highlight the complexity of CRL functions and substrate interactions in transcriptional reprogramming during establishment of immunity.

Beyond E3 ligases the proteasome itself may also play important roles in the regulation of gene expression programmes. In yeast and human cells the proteasome has been found to physically associate with the chromatin and regulate the expression of thousands of genes (Collins and Tansey, 2006; Geng *et al.*, 2012). Our understanding of how proteasomes are recruited to the chromatin sites where they are most needed is still in its infancy but may be dependent on both E3 ligases and their substrates. In this respect it is interesting to note that many E3 ligases interact with the 19S proteasome particle, suggesting they might directly hand over ubiquitinated substrates for degradation (Schmidt *et al.*, 2005). Moreover, the proteasome may have resident E3 ligases that further modify substrates before their degradation (Crosas *et al.*, 2006; Schmidt *et al.*, 2005), indicating further signalling complexity is achieved at the proteasome itself. Thus, to gain full appreciation of how the UPS controls transcriptional reprogramming in plant immunity, the future challenge is to uncover post-translational regulation and substrate repertoires of E3 ligases and the proteasome itself across different interconnected immune signalling pathways.

Acknowledgements

S.H.S was supported by a Royal Society University Research Fellowship (UF140600), the Biotechnology and Biological Sciences Research Council (BBSRC) grant no. BB/L006219/1 and by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme Grant No. 678511. E.H.G.A. was supported by an

463 Industrial CASE studentship (BB/M503216/1) from the BBSRC co-funded by Bayer Crop
464 Science AG.
465

References

- Boyle P, Le Su E, Rochon A, Shearer HL, Murmu J, Chu JY, Fobert PR, Despres C.** 2009. The BTB/POZ domain of the Arabidopsis disease resistance protein NPR1 interacts with the repression domain of TGA2 to negate its function. *The Plant Cell* **21**, 3700-3713.
- Canet JV, Dobon A, Fajmonova J, Tornero P.** 2012. The *BLADE-ON-PETIOLE* genes of Arabidopsis are essential for resistance induced by methyl jasmonate. *BMC Plant Biology* **12**, 199.
- Canet JV, Dobon A, Roig A, Tornero P.** 2010. Structure-function analysis of *npr1* alleles in Arabidopsis reveals a role for its paralogs in the perception of salicylic acid. *Plant Cell & Environment* **33**, 1911-1922.
- Canning P, Cooper CDO, Krojer T, Murray JW, Pike AC, Chaikuad A, Keates T, Thangaratnarajah C, Hojzan V, Ayinampudi V, Marsden BD, Gileadi O, Knapp S, von Delft F, Bullock AN.** 2013. Structural basis for Cul3 protein assembly with the BTB-Kelch family of E3 ubiquitin ligases. *Journal of Biological Chemistry* **288**, 7803-7814.
- Cao H, Bowling SA, Gordon AS, Dong X.** 1994. Characterization of an Arabidopsis mutant that is nonresponsive to inducers of systemic acquired resistance. *The Plant Cell* **6**, 1583-1592.
- Cao H, Glazebrook J, Clarke JD, Volko S, Dong X.** 1997. The Arabidopsis NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell* **88**, 57-63.
- Chen H, Chen J, Li M, Chang M, Xu K, Shang Z, Zhao Y, Palmer I, Zhang Y, McGill J, Alfano JR, Nishimura MT, Liu F, Fu ZQ.** 2017. A bacterial type III effector targets the master regulator of salicylic acid signaling, NPR1, to subvert plant immunity. *Cell Host & Microbe* **22**, 777-788 e777.
- Chern M-S, Fitzgerald HA, Yadav RC, Canlas PE, Dong X, Ronald PC.** 2001. Evidence for a disease-resistance pathway in rice similar to the *NPR1*-mediated signaling pathway in Arabidopsis. *Plant Journal* **27**, 101-113.
- Chern M, Bai W, Ruan D, Oh T, Chen X, Ronald PC.** 2014. Interaction specificity and coexpression of rice NPR1 homologs 1 and 3 (NH1 and NH3), TGA transcription factors and Negative Regulator of Resistance (NRR) proteins. *BMC Genomics* **15**, 461.
- Chern M, Fitzgerald HA, Canlas PE, Navarre DA, Ronald PC.** 2005. Overexpression of a rice NPR1 homolog leads to constitutive activation of defense response and hypersensitivity to light. *Molecular Plant-Microbe Interactions* **18**, 511-520.
- Collins GA, Tansey WP.** 2006. The proteasome: a utility tool for transcription? *Current Opinion in Genetics & Development* **16**, 197-202.
- Crosas B, Hanna J, Kirkpatrick DS, Zhang DP, Tone Y, Hathaway NA, Buecker C, Leggett DS, Schmidt M, King RW, Gygi SP, Finley D.** 2006. Ubiquitin chains are remodeled at the proteasome by opposing ubiquitin ligase and deubiquitinating activities. *Cell* **127**, 1401-1413.
- de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, Mansfield JW, Rodriguez Egea P, Bogre L, Grant M.** 2007. *Pseudomonas syringae* pv. *tomato* hijacks the Arabidopsis abscisic acid signalling pathway to cause disease. *Embo Journal* **26**, 1434-1443.
- Dharmasiri N, Dharmasiri S, Estelle M.** 2005. The F-box protein TIR1 is an auxin receptor. *Nature* **435**, 441-445.
- Delaney TP, Friedrich L, Ryals JA.** 1995. Arabidopsis signal transduction mutant defective in chemically and biologically induced disease resistance. *Proceedings of the National Academy of Sciences USA* **92**, 6602-6606.
- Després C, DeLong C, Glaze S, Liu E, Fobert PR.** 2000. The Arabidopsis NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *The Plant Cell* **12**, 279-290.

Ding Y, Dommel M, Mou Z. 2016. Absciscic Acid Promotes Proteasome-Mediated Degradation of the Transcription Coactivator NPR1 in *Arabidopsis thaliana*. *The Plant Journal*.

Du L, Ali GS, Simons KA, Hou J, Yang T, Reddy ASN, Poovaiah BW. 2009. Ca^{2+} /calmodulin regulates salicylic-acid-mediated plant immunity. *Nature* **457**, 1154-1158.

Duplan V, Rivas S. 2014. E3 ubiquitin-ligases and their target proteins during the regulation of plant innate immunity. *Frontiers in Plant Science* **5**, 42.

Fu ZQ, Yan S, Saleh A, Wang W, Ruble J, Oka N, Mohan R, Spoel SH, Tada Y, Zheng N, Dong X. 2012. NPR3 and NPR4 are receptors for the immune signal salicylic acid in plants. *Nature* **486**, 228-232.

Furniss JJ, Spoel SH. 2015. CULLIN-RING ubiquitin ligases in salicylic acid-mediated plant immune signaling. *Frontiers in Plant Science* **6**, 154.

Gagne JM, Downes BP, Shiu SH, Durski AM, Vierstra RD. 2002. The F-box subunit of the SCF E3 complex is encoded by a diverse superfamily of genes in *Arabidopsis*. *Proceedings of the National Academy of Sciences USA* **99**, 11519–11524.

Geng F, Wenzel S, Tansey WP. 2012. Ubiquitin and proteasomes in transcription. *Annual Review of Biochemistry* **81**, 177-201.

Glazebrook J, Rogers EE, Ausubel FM. 1996. Isolation of *Arabidopsis* mutants with enhanced disease susceptibility by direct screening. *Genetics* **143**, 973-982.

Goossens J, Fernandez-Calvo P, Schweizer F, Goossens A. 2016. Jasmonates: signal transduction components and their roles in environmental stress responses. *Plant Molecular Biology* **91**, 673-689.

Hepworth SR, Zhang Y, McKim S, Li X, Haughn GW. 2005. BLADE-ON-PETIOLE-dependent signaling controls leaf and floral patterning in *Arabidopsis*. *The Plant Cell* **17**, 1434-1448.

Kelley DR, Estelle M. 2012. Ubiquitin-mediated control of plant hormone signaling. *Plant Physiology* **160**, 47-55.

Kepinski S, Leyser O. 2005. The *Arabidopsis* F-box protein TIR1 is an auxin receptor. *Nature* **435**, 446–451.

Kinkema M, Fan W, Dong X. 2000. Nuclear localization of NPR1 is required for activation of *PR* gene expression. *The Plant Cell* **12**, 2339-2350.

Komander D, Rape M. 2012. The ubiquitin code. *Annual Review of Biochemistry* **81**, 203-229.

Lee HJ, Park YJ, Seo PJ, Kim JH, Sim HJ, Kim SG, Park CM. 2015. Systemic Immunity Requires SnRK2.8-Mediated Nuclear Import of NPR1 in *Arabidopsis*. *The Plant Cell* **27**, 3425-3438.

Liu G, Holub EB, Alonso JM, Ecker JR, Fobert PR. 2005. An *Arabidopsis* *NPR1*-like gene, *NPR4*, is required for disease resistance. *Plant Journal* **41**, 304-318.

Liu L, Sonbol FM, Huot B, Gu Y, Withers J, Mwimba M, Yao J, He SY, Dong X. 2016. Salicylic acid receptors activate jasmonic acid signalling through a non-canonical pathway to promote effector-triggered immunity. *Nature Communications* **7**, 13099.

Liu Q, Ning Y, Zhang Y, Yu N, Zhao C, Zhan X, Wu W, Chen D, Wei X, Wang GL, Cheng S, Cao L. 2017. OsCUL3a Negatively Regulates Cell Death and Immunity by Degrading OsNPR1 in Rice. *The Plant Cell* **29**, 345-359.

Marino D, Peeters N, Rivas S. 2012. Ubiquitination during plant immune signaling. *Plant Physiology* **160**, 15-27.

Matsushita A, Inoue H, Goto S, Nakayama A, Sugano S, Hayashi N, Takatsuji H. 2013. Nuclear ubiquitin proteasome degradation affects WRKY45 function in the rice defense program. *The Plant Journal* **73**, 302-313.

Mazzucotelli E, Belloni S, Marone D, De Leonardis A, Guerra D, Di Fonzo N, Cattivelli L, Mastrangelo A. 2006. The e3 ubiquitin ligase gene family in plants: regulation by degradation. *Current Genomics* **7**, 509-522.

McMahon M, Thomas N, Itoh K, Yamamoto M, Hayes JD. 2006. Dimerization of substrate adaptors can facilitate cullin-mediated ubiquitylation of proteins by a "tethering" mechanism:

a two-site interaction model for the Nrf2-Keap1 complex. *Journal of Biological Chemistry* **281**, 24756-24768.

Miao Y, Zentgraf U. 2010. A HECT E3 ubiquitin ligase negatively regulates Arabidopsis leaf senescence through degradation of the transcription factor WRKY53. *The Plant Journal* **63**, 179-188.

Mohr PG, Cahill DM. 2007. Suppression by ABA of salicylic acid and lignin accumulation and the expression of multiple genes, in *Arabidopsis* infected with *Pseudomonas syringae* pv. *tomato*. *Functional & Integrative Genomics* **7**, 181-191.

Mou Z, Fan W, Dong X. 2003. Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* **113**, 935-944.

Nakayama A, Fukushima S, Goto S, Matsushita A, Shimono M, Sugano S, Jiang CJ, Akagi A, Yamazaki M, Inoue H, Takatsuji H. 2013. Genome-wide identification of WRKY45-regulated genes that mediate benzothiadiazole-induced defense responses in rice. *BMC Plant Biology* **13**, 150.

Ni W, Xu SL, Tepperman JM, Stanley DJ, Maltby DA, Gross JD, Burlingame AL, Wang ZY, Quail PH. 2014. A mutually assured destruction mechanism attenuates light signaling in *Arabidopsis*. *Science* **344**, 1160-1164.

Petroski MD, Deshaies RJ. 2005. Function and regulation of cullin-RING ubiquitin ligases. *Nature Reviews Molecular Cell Biology* **6**, 9-20.

Pickart CM, Cohen RE. 2004. Proteasomes and their kin: proteases in the machine age. *Nature Reviews Molecular Cell Biology* **5**, 177-187.

Pontier D, Privat I, Trifa Y, Zhou JM, Klessig DF, Lam E. 2002. Differential regulation of TGA transcription factors by post-transcriptional control. *The Plant Journal* **32**, 641-653.

Rochon A, Boyle P, Wignes T, Fobert PR, Després C. 2006. The coactivator function of *Arabidopsis* NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal cysteines. *The Plant Cell* **18**, 3670-3685.

Ryals J, Weymann K, Lawton K, Friedrich L, Ellis D, Steiner HY, Johnson J, Delaney TP, Jesse T, Vos P, Uknes S. 1997. The Arabidopsis NIM1 protein shows homology to the mammalian transcription factor inhibitor IκB. *The Plant Cell* **9**, 425-439.

Saleh A, Withers J, Mohan R, Marqués J, Gu Y, Yan S, Zavaliev R, Nomoto M, Tada Y, Dong X. 2015. Posttranslational Modifications of the Master Transcriptional Regulator NPR1 Enable Dynamic but Tight Control of Plant Immune Responses. *Cell Host & Microbe* **18**, 169-182.

Salghetti SE, Muratani M, Wijnen H, Futcher B, Tansey WP. 2000. Functional overlap of sequences that activate transcription and signal ubiquitin-mediated proteolysis. *Proceedings of the National Academy of Sciences USA* **97**, 3118-3123.

Santner A, Estelle M. 2009. Recent advances and emerging trends in plant hormone signalling. *Nature* **459**, 1071-1078.

Schmidt M, Hanna J, Elsasser S, Finley D. 2005. Proteasome-associated proteins: regulation of a proteolytic machine. *Biol Chem* **386**, 725-737.

Seibert V, Prohl C, Schoultz I, Rhee E, Lopez R, Abderazzaq K, Zhou C, Wolf DA. 2002. Combinatorial diversity of fission yeast SCF ubiquitin ligases by homo- and heterooligomeric assemblies of the F-box proteins Pop1p and Pop2p. *BMC Biochemistry* **3**, 22.

Shah J, Tsui F, Klessig DF. 1997. Characterization of a salicylic acid-insensitive mutant (sai1) of *Arabidopsis thaliana*, identified in a selective screen utilizing the SA-inducible expression of the tms2 gene. *Molecular Plant-Microbe Interactions* **10**, 69-78.

Shi Z, Maximova S, Liu Y, Verica J, Guiltinan MJ. 2013. The salicylic acid receptor NPR3 is a negative regulator of the transcriptional defense response during early flower development in *Arabidopsis*. *Molecular Plant* **6**, 802-816.

Shimono M, Koga H, Akagi A, Hayashi N, Goto S, Sawada M, Kurihara T, Matsushita A, Sugano S, Jiang CJ, Kaku H, Inoue H, Takatsuji H. 2012. Rice WRKY45 plays important roles in fungal and bacterial disease resistance. *Molecular Plant Pathology* **13**, 83-94.

Shimono M, Sugano S, Nakayama A, Jiang CJ, Ono K, Toki S, Takatsuji H. 2007. Rice WRKY45 plays a crucial role in benzothiadiazole-inducible blast resistance. *The Plant Cell* **19**, 2064-2076.

Skelly MJ, Frungillo L, Spoel SH. 2016. Transcriptional regulation by complex interplay between post-translational modifications. *Current Opinion in Plant Biology* **33**, 126-132.

Smalle J, Vierstra RD. 2004. The ubiquitin 26S proteasome proteolytic pathway. *Annual Review of Plant Biology* **55**, 555-590.

Spoel SH, Dong X. 2012. How do plants achieve immunity? Defence without specialized immune cells. *Nature Reviews Immunology* **12**, 89-100.

Spoel SH, Johnson JS, Dong X. 2007. Regulation of tradeoffs between plant defenses against pathogens with different lifestyles. *Proceedings of the National Academy of Sciences USA* **104**, 18842-18847.

Spoel SH, Koornneef A, Claessens SMC, Korzelius JP, Van Pelt JA, Mueller MJ, Buchala AJ, Métraux J-P, Brown R, Kazan K, Van Loon LC, Dong X, Pieterse CMJ. 2003. NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *The Plant Cell* **15**, 760-770.

Spoel SH, Mou Z, Tada Y, Spivey NW, Genschik P, Dong X. 2009. Proteasome-mediated turnover of the transcription coactivator NPR1 plays dual roles in regulating plant immunity. *Cell* **137**, 860-872.

Stogios PJ, Chen L, Prive GG. 2007. Crystal structure of the BTB domain from the LRF/ZBTB7 transcriptional regulator. *Protein Science* **16**, 336-342.

Stogios PJ, Downs GS, Jauhal JJS, Nandra SK, Prive GG. 2005. Sequence and structural analysis of BTB domain proteins. *Genome Biol* **6**, R82.

Sugano S, Jiang CJ, Miyazawa SI, Masumoto C, Yazawa K, Hayashi N, Shimono M, Nakayama A, Miyao M, Takatsuji H. 2010. Role of OsNPR1 in rice defense program as revealed by genome-wide expression analysis. *Plant Molecular Biology* **74**, 549-562.

Tada Y, Spoel SH, Pajerowska-Mukhtar K, Mou Z, Song J, Wang C, Zuo J, Dong X. 2008. Plant immunity requires conformational changes of NPR1 via S-nitrosylation and thioredoxins. *Science* **321**, 952-956.

Thrower JS, Hoffman L, Rechsteiner M, Pickart CM. 2000. Recognition of the polyubiquitin proteolytic signal. *Embo Journal* **19**, 94-102.

Trujillo M, Shirasu K. 2010. Ubiquitination in plant immunity. *Current Opinion in Plant Biology* **13**, 402-408.

Üstün S, Bartetzko V, Bornke F. 2013. The *Xanthomonas campestris* type III effector XopJ targets the host cell proteasome to suppress salicylic-acid mediated plant defence. *PLoS Pathogens* **9**, e1003427.

Üstün S, Bornke F. 2015. The *Xanthomonas campestris* type III effector XopJ proteolytically degrades proteasome subunit RPT6. *Plant Physiology* **168**, 107-119.

Üstün S, Sheikh A, Gimenez-Ibanez S, Jones A, Ntoukakis V, Bornke F. 2016. The proteasome acts as a hub for plant immunity and is targeted by *Pseudomonas* type-III effectors. *Plant Physiology* **172**, 1941-1958.

Van der Does D, Leon-Reyes A, Koornneef A, Van Verk MC, Rodenburg N, Pauwels L, Goossens A, Korbes AP, Memelink J, Ritsema T, Van Wees SCM, Pieterse CMJ. 2013. Salicylic acid suppresses jasmonic acid signaling downstream of SCF^{COI1}-JAZ by targeting GCC promoter motifs via transcription factor ORA59. *The Plant Cell* **25**, 744-761.

Vierstra RD. 2009. The ubiquitin-26S proteasome system at the nexus of plant biology. *Nature Reviews Molecular Cell Biology* **10**, 385-397.

Wang D, Amornsiripanitch N, Dong X. 2006. A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. *PLoS Pathogens* **2**, e123.

Weigel RR, Bauscher C, Pfitzner AJ, Pfitzner UM. 2001. NIMIN-1, NIMIN-2 and NIMIN-3, members of a novel family of proteins from Arabidopsis that interact with NPR1/NIM1, a key regulator of systemic acquired resistance in plants. *Plant Molecular Biology* **46**, 143-160.

Yasuda M, Ishikawa A, Jikumaru Y, Seki M, Umezawa T, Asami T, Maruyama-Nakashita A, Kudo T, Shinozaki K, Yoshida S, Nakashita H. 2008. Antagonistic interaction between systemic acquired resistance and the abscisic acid-mediated abiotic stress response in Arabidopsis. *The Plant Cell* **20**, 1678-1692.

Yu Y, Xu W, Wang J, Wang L, Yao W, Yang Y, Xu Y, Ma F, Du Y, Wang Y. 2013. The Chinese wild grapevine (*Vitis pseudoreticulata*) E3 ubiquitin ligase *Erysiphe necator*-induced RING finger protein 1 (EIRP1) activates plant defense responses by inducing proteolysis of the VpWRKY11 transcription factor. *New Phytologist* **200**, 834-846.

Yuan Y, Zhong S, Li Q, Zhu Z, Lou Y, Wang L, Wang J, Wang M, Yang D, He Z. 2007. Functional analysis of rice NPR1-like genes reveals that OsNPR1/NH1 is the rice orthologue conferring disease resistance with enhanced herbivore susceptibility. *Plant Biotechnology Journal* **5**, 313-324.

Zhang L, Du L, Shen C, Yang Y, Poovaiah BW. 2014. Regulation of plant immunity through ubiquitin-mediated modulation of Ca(2+) -calmodulin-AtSR1/CAMTA3 signaling. *The Plant Journal* **78**, 269-281.

Zhang L, Zhang F, Melotto M, Yao J, He SY. 2017. Jasmonate signaling and manipulation by pathogens and insects. *Journal of Experimental Botany* **68**, 1371-1385.

Zhang Y, Cheng YT, Qu N, Zhao Q, Bi D, Li X. 2006. Negative regulation of defense responses in Arabidopsis by two *NPR1* paralogs. *The Plant Journal* **48**, 647-656.

Zhang Y, Fan W, Kinkema M, Li X, Dong X. 1999. Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the *PR-1* gene. *Proceedings of the National Academy of Sciences USA* **96**, 6523-6528.

Zheng N, Shabek N. 2017. Ubiquitin Ligases: Structure, Function, and Regulation. *Annual Review Biochemistry* **86**, 129-157.

Zhou JM, Trifa Y, Silva H, Pontier D, Lam E, Shah J, Klessig DF. 2000. NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the *PR-1* gene required for induction by salicylic acid. *Molecular Plant-Microbe Interactions* **13**, 191-202.

Zhuang M, Calabrese MF, Liu J, Waddell MB, Nourse A, Hammel M, Miller DJ, Walden H, Duda DM, Seyedin SN, Hoggard T, Harper JW, White KP, Schulman BA. 2009. Structures of SPOP-substrate complexes: insights into molecular architectures of BTB-Cul3 ubiquitin ligases. *Molecular Cell* **36**, 39-50.

Zimmerman ES, Schulman BA, Zheng N. 2010. Structural assembly of cullin-RING ubiquitin ligase complexes. *Current Opinion in Structural Biology* **20**, 714-721.






Figure 1. CRL3-mediated transcriptional reprogramming in Arabidopsis and rice

In Arabidopsis (left panel) stability of the transcriptional coactivator NPR1 is controlled by different CRL3 ligases. In unchallenged cells NPR1 is polyubiquitinated (grey circles) by CRL3^{NPR4} ligases to prevent autoimmunity (*i.e.* immunosuppression). Upon pathogen challenge SA induces the SUMOylation (S, blue circles) and phosphorylation (P, green circles) of NPR1, allowing transcriptionally active NPR1 to undergo a transcription-coupled proteolysis cycle that stimulates the expression of immune genes. In rice (right panel) immunosuppression is accomplished by CRL3-mediated ubiquitination and degradation of OsNPR1 as well as by degradation of OsWRKY45 which may also involve a CRL3 ligase. Upon pathogen challenge OsNPR1 activates immune genes but also suppresses genes involved in growth and development. Additionally, pathogen challenge activates OsWRKY45, resulting in its phosphorylation and subsequent degradation in a proteolysis-coupled transcription cycle that is hypothesised to involve a CRL3 ligase. Note that the transcriptionally competent or active state of all transcription (co)regulators is represented by a shaded box.

Arabidopsis

Rice



Immunosuppression

Transcriptionally active

Proteolysis-coupled transcription cycle